

Figure 31.15 The German-born American physicist Maria Goeppert Mayer (1906–1972) shared the 1963 Nobel Prize in physics with J. Jensen for the creation of the nuclear shell model. This successful nuclear model has nucleons filling shells analogous to electron shells in atoms. It was inspired by patterns observed in nuclear properties. (credit: Nobel Foundation via Wikimedia Commons)

## **31.4 Nuclear Decay and Conservation Laws**

Nuclear **decay** has provided an amazing window into the realm of the very small. Nuclear decay gave the first indication of the connection between mass and energy, and it revealed the existence of two of the four basic forces in nature. In this section, we explore the major modes of nuclear decay; and, like those who first explored them, we will discover evidence of previously unknown particles and conservation laws.

Some nuclides are stable, apparently living forever. Unstable nuclides decay (that is, they are radioactive), eventually producing a stable nuclide after many decays. We call the original nuclide the **parent** and its decay products the **daughters**. Some radioactive nuclides decay in a single step to a stable nucleus. For example, <sup>60</sup>Co is unstable and decays directly to <sup>60</sup>Ni, which is stable. Others, such as <sup>238</sup>U, decay to another unstable nuclide, resulting in a **decay series** in which each subsequent nuclide decays until a stable nuclide is finally produced. The decay series that starts from <sup>238</sup>U is of particular interest, since it produces the radioactive isotopes <sup>226</sup>Ra and <sup>210</sup>Po, which the Curies first discovered (see Figure 31.16). Radon gas is also produced (<sup>222</sup>Rn in the series), an increasingly recognized naturally occurring hazard. Since radon is a noble gas, it emanates from materials, such as soil, containing even trace amounts of <sup>238</sup>U and can be inhaled. The decay of radon and its daughters produces internal damage. The <sup>238</sup>U decay series ends with <sup>206</sup>Pb, a stable isotope of lead.



**Figure 31.16** The decay series produced by <sup>238</sup> U, the most common uranium isotope. Nuclides are graphed in the same manner as in the chart of nuclides. The type of decay for each member of the series is shown, as well as the half-lives. Note that some nuclides decay by more than one mode. You can see why radium and polonium are found in uranium ore. A stable isotope of lead is the end product of the series.

Note that the daughters of  $\alpha$  decay shown in Figure 31.16 always have two fewer protons and two fewer neutrons than the parent. This seems reasonable, since we know that  $\alpha$  decay is the emission of a <sup>4</sup>He nucleus, which has two protons and two neutrons. The daughters of  $\beta$  decay have one less neutron and one more proton than their parent. Beta decay is a little more subtle, as we shall see. No  $\gamma$  decays are shown in the figure, because they do not produce a daughter that differs from the parent.

## **Alpha Decay**

In **alpha decay**, a <sup>4</sup>He nucleus simply breaks away from the parent nucleus, leaving a daughter with two fewer protons and two fewer neutrons than the parent (see Figure 31.17). One example of  $\alpha$  decay is shown in Figure 31.16 for <sup>238</sup>U. Another nuclide that undergoes  $\alpha$  decay is <sup>239</sup>Pu. The decay equations for these two nuclides are

$$^{238}\text{U} \rightarrow ^{234}\text{Th}_{92}^{234} + {}^{4}\text{He}$$
 31.13

and

$$^{239}$$
Pu  $\rightarrow ^{235}$ U + <sup>4</sup>He. 31.14



Figure 31.17 Alpha decay is the separation of a <sup>4</sup>He nucleus from the parent. The daughter nucleus has two fewer protons and two fewer neutrons than the parent. Alpha decay occurs spontaneously only if the daughter and <sup>4</sup>He nucleus have less total mass than the parent.

If you examine the periodic table of the elements, you will find that Th has Z = 90, two fewer than U, which has Z = 92. Similarly, in the second **decay equation**, we see that U has two fewer protons than Pu, which has Z = 94. The general rule for  $\alpha$ decay is best written in the format  ${}^{A}_{A}X_{N}$ . If a certain nuclide is known to  $\alpha$  decay (generally this information must be looked up in a table of isotopes, such as in Appendix B), its  $\alpha$  decay equation is

$${}^{A}_{Z}X_{N} \rightarrow {}^{A-4}_{Z-2}Y_{N-2} + {}^{4}_{2}\text{He}_{2} \quad (\alpha \text{ decay})$$
 31.15

where Y is the nuclide that has two fewer protons than X, such as Th having two fewer than U. So if you were told that  $^{239}$ Pu  $\alpha$ decays and were asked to write the complete decay equation, you would first look up which element has two fewer protons (an atomic number two lower) and find that this is uranium. Then since four nucleons have broken away from the original 239, its atomic mass would be 235.

It is instructive to examine conservation laws related to  $\alpha$  decay. You can see from the equation  ${}^{A}_{Z}X_{N} \rightarrow {}^{A-4}_{Z-2}Y_{N-2} + {}^{4}_{2}\text{He}_{2}$ that total charge is conserved. Linear and angular momentum are conserved, too. Although conserved angular momentum is not of great consequence in this type of decay, conservation of linear momentum has interesting consequences. If the nucleus is at rest when it decays, its momentum is zero. In that case, the fragments must fly in opposite directions with equal-magnitude momenta so that total momentum remains zero. This results in the  $\alpha$  particle carrying away most of the energy, as a bullet from a heavy rifle carries away most of the energy of the powder burned to shoot it. Total mass-energy is also conserved: the energy produced in the decay comes from conversion of a fraction of the original mass. As discussed in Atomic Physics, the general relationship is

$$E = (\Delta m)c^2.$$
 31.16

Here, E is the **nuclear reaction energy** (the reaction can be nuclear decay or any other reaction), and  $\Delta m$  is the difference in mass between initial and final products. When the final products have less total mass,  $\Delta m$  is positive, and the reaction releases energy (is exothermic). When the products have greater total mass, the reaction is endothermic ( $\Delta m$  is negative) and must be induced with an energy input. For  $\alpha$  decay to be spontaneous, the decay products must have smaller mass than the parent.

# **EXAMPLE 31.2**

## **Alpha Decay Energy Found from Nuclear Masses**

Find the energy emitted in the  $\alpha$  decay of <sup>239</sup>Pu.

### Strategy

Nuclear reaction energy, such as released in  $\alpha$  decay, can be found using the equation  $E = (\Delta m)c^2$ . We must first find  $\Delta m$ , the difference in mass between the parent nucleus and the products of the decay. This is easily done using masses given in Appendix Α.

### Solution

The decay equation was given earlier for  $^{239}$ Pu ; it is

 $^{239}$ Pu  $\rightarrow ^{235}$ U +  $^{4}$ He

31.17

Thus the pertinent masses are those of  $^{239}$  Pu,  $^{235}$  U, and the  $\alpha$  particle or  $^{4}$  He, all of which are listed in Appendix A. The initial mass was  $m(^{239}\text{Pu}) = 239.052157 \text{ u}$ . The final mass is the sum

 $m(^{235}\text{U}) + m(^{4}\text{He}) = 235.043924 \text{ u} + 4.002602 \text{ u} = 239.046526 \text{ u}$ . Thus,

$$\Delta m = m(^{239}\text{Pu}) - [m(^{235}\text{U}) + m(^{4}\text{He})]$$
  
= 239.052157 u - 239.046526 u  
= 0.0005631 u.

Now we can find *E* by entering  $\Delta m$  into the equation:

$$E = (\Delta m)c^2 = (0.005631 \text{ u})c^2.$$

We know 1  $u = 931.5 \text{ MeV}/c^2$ , and so

$$E = (0.005631)(931.5 \text{ MeV}/c^2)(c^2) = 5.25 \text{ MeV}.$$

#### Discussion

The energy released in this  $\alpha$  decay is in the MeV range, about  $10^6$  times as great as typical chemical reaction energies, consistent with many previous discussions. Most of this energy becomes kinetic energy of the  $\alpha$  particle (or <sup>4</sup>He nucleus), which moves away at high speed. The energy carried away by the recoil of the <sup>235</sup>U nucleus is much smaller in order to conserve momentum. The <sup>235</sup>U nucleus can be left in an excited state to later emit photons ( $\gamma$  rays). This decay is spontaneous and releases energy, because the products have less mass than the parent nucleus. The question of why the products have less mass will be discussed in <u>Binding Energy</u>. Note that the masses given in <u>Appendix A</u> are atomic masses of neutral atoms, including their electrons. The mass of the electrons is the same before and after  $\alpha$  decay, and so their masses subtract out when finding  $\Delta m$ . In this case, there are 94 electrons before and after the decay.

## **Beta Decay**

There are actually *three* types of **beta decay**. The first discovered was "ordinary" beta decay and is called  $\beta^-$  decay or electron emission. The symbol  $\beta^-$  represents *an electron emitted in nuclear beta decay*. Cobalt-60 is a nuclide that  $\beta^-$  decays in the following manner:

$${}^{60}\text{Co} \rightarrow {}^{60}\text{Ni} + \beta^- + \text{neutrino.}$$

The **neutrino** is a particle emitted in beta decay that was unanticipated and is of fundamental importance. The neutrino was not even proposed in theory until more than 20 years after beta decay was known to involve electron emissions. Neutrinos are so difficult to detect that the first direct evidence of them was not obtained until 1953. Neutrinos are nearly massless, have no charge, and do not interact with nucleons via the strong nuclear force. Traveling approximately at the speed of light, they have little time to affect any nucleus they encounter. This is, owing to the fact that they have no charge (and they are not EM waves), they do not interact through the EM force. They do interact via the relatively weak and very short range weak nuclear force. Consequently, neutrinos escape almost any detector and penetrate almost any shielding. However, neutrinos do carry energy, angular momentum (they are fermions with half-integral spin), and linear momentum away from a beta decay. When accurate measurements of beta decay were made, it became apparent that energy, angular momentum, and linear momentum were not three conservation laws were being violated. Wolfgang Pauli made a formal proposal for the existence of neutrinos in 1930. The Italian-born American physicist Enrico Fermi (1901–1954) gave neutrinos their name, meaning little neutral ones, when he developed a sophisticated theory of beta decay (see Figure 31.18). Part of Fermi's theory was the identification of the weak nuclear force as being distinct from the strong nuclear force and in fact responsible for beta decay.

31.19

31.20

31.21



Figure 31.18 Enrico Fermi was nearly unique among 20th-century physicists—he made significant contributions both as an experimentalist and a theorist. His many contributions to theoretical physics included the identification of the weak nuclear force. The fermi (fm) is named after him, as are an entire class of subatomic particles (fermions), an element (Fermium), and a major research laboratory (Fermilab). His experimental work included studies of radioactivity, for which he won the 1938 Nobel Prize in physics, and creation of the first nuclear chain reaction. (credit: United States Department of Energy, Office of Public Affairs)

The neutrino also reveals a new conservation law. There are various families of particles, one of which is the electron family. We propose that the number of members of the electron family is constant in any process or any closed system. In our example of beta decay, there are no members of the electron family present before the decay, but after, there is an electron and a neutrino. So electrons are given an electron family number of +1. The neutrino in  $\beta^-$  decay is an **electron's antineutrino**, given the symbol  $\overline{\nu}_e$ , where  $\nu$  is the Greek letter nu, and the subscript e means this neutrino is related to the electron. The bar indicates this is a particle of **antimatter**. (All particles have antimatter counterparts that are nearly identical except that they have the opposite charge. Antimatter is almost entirely absent on Earth, but it is found in nuclear decay and other nuclear and particle reactions as well as in outer space.) The electron's antineutrino  $\overline{\nu}_e$ , being antimatter, has an electron family number of -1. The total is zero, before and after the decay. The new conservation law, obeyed in all circumstances, states that the *total electron family number is constant*. An electron cannot be created without also creating an antimatter family member. This law is analogous to the conservation of charge in a situation where total charge is originally zero, and equal amounts of positive and negative charge must be created in a reaction to keep the total zero.

If a nuclide  ${}^A_Z X_N$  is known to  $\beta^-$  decay, then its  $\beta^-$  decay equation is

where Y is the nuclide having one more proton than X (see Figure 31.19). So if you know that a certain nuclide  $\beta^-$  decays, you can find the daughter nucleus by first looking up Z for the parent and then determining which element has atomic number Z + 1. In the example of the  $\beta^-$  decay of <sup>60</sup>Co given earlier, we see that Z = 27 for Co and Z = 28 is Ni. It is as if one of the neutrons in the parent nucleus decays into a proton, electron, and neutrino. In fact, neutrons outside of nuclei do just that—they live only an average of a few minutes and  $\beta^-$  decay in the following manner:



Figure 31.19 In  $\beta^-$  decay, the parent nucleus emits an electron and an antineutrino. The daughter nucleus has one more proton and one less neutron than its parent. Neutrinos interact so weakly that they are almost never directly observed, but they play a fundamental role in

### particle physics.

We see that charge is conserved in  $\beta^-$  decay, since the total charge is Z before and after the decay. For example, in <sup>60</sup>Co decay, total charge is 27 before decay, since cobalt has Z = 27. After decay, the daughter nucleus is Ni, which has Z = 28, and there is an electron, so that the total charge is also 28 + (-1) or 27. Angular momentum is conserved, but not obviously (you have to examine the spins and angular momenta of the final products in detail to verify this). Linear momentum is also conserved, again imparting most of the decay energy to the electron and the antineutrino, since they are of low and zero mass, respectively. Another new conservation law is obeyed here and elsewhere in nature. *The total number of nucleons A is conserved*. In <sup>60</sup>Co decay, for example, there are 60 nucleons before and after the decay. Note that total *A* is also conserved in  $\alpha$  decay. Also note that the total number of protons changes, as does the total number of neutrons, so that total *X* and total *N* are *not* conserved in  $\beta^-$  decay, as they are in  $\alpha$  decay. Energy released in  $\beta^-$  decay can be calculated given the masses of the parent and products.

# EXAMPLE 31.3

## $\beta^-$ Decay Energy from Masses

Find the energy emitted in the  $\beta^-$  decay of  ${}^{60}$ Co.

### **Strategy and Concept**

As in the preceding example, we must first find  $\Delta m$ , the difference in mass between the parent nucleus and the products of the decay, using masses given in <u>Appendix A</u>. Then the emitted energy is calculated as before, using  $E = (\Delta m)c^2$ . The initial mass is just that of the parent nucleus, and the final mass is that of the daughter nucleus and the electron created in the decay. The neutrino is massless, or nearly so. However, since the masses given in <u>Appendix A</u> are for neutral atoms, the daughter nucleus has one more electron than the parent, and so the extra electron mass that corresponds to the  $\beta^-$  is included in the atomic mass of Ni. Thus,

$$\Delta m = m({}^{60}\text{Co}) - m({}^{60}\text{Ni}).$$
31.24

#### Solution

The  $\beta^-$  decay equation for <sup>60</sup>Co is

$${}^{60}_{27}\text{Co}_{33} \rightarrow {}^{60}_{28}\text{Ni}_{32} + \beta^- + \overline{\nu}_e.$$
 31.25

As noticed,

$$\Delta m = m({}^{60}\text{Co}) - m({}^{60}\text{Ni}). \qquad 31.26$$

Entering the masses found in Appendix A gives

$$\Delta m = 59.933820 \text{ u} - 59.930789 \text{ u} = 0.003031 \text{ u}.$$

Thus,

$$E = (\Delta m)c^2 = (0.003031 \text{ u})c^2$$
.

Using 1 u = 931.5 MeV/ $c^2$ , we obtain

$$E = (0.003031)(931.5 \text{ MeV}/c^2)(c^2) = 2.82 \text{ MeV}.$$
 31.29

### **Discussion and Implications**

Perhaps the most difficult thing about this example is convincing yourself that the  $\beta^-$  mass is included in the atomic mass of  ${}^{60}$ Ni. Beyond that are other implications. Again the decay energy is in the MeV range. This energy is shared by all of the products of the decay. In many  ${}^{60}$ Co decays, the daughter nucleus  ${}^{60}$ Ni is left in an excited state and emits photons ( $\gamma$  rays). Most of the remaining energy goes to the electron and neutrino, since the recoil kinetic energy of the daughter nucleus is small. One final note: the electron emitted in  $\beta^-$  decay is created in the nucleus at the time of decay.

The second type of beta decay is less common than the first. It is  $\beta^+$  decay. Certain nuclides decay by the emission of a *positive* electron. This is **antielectron** or **positron decay** (see Figure 31.20).



Figure 31.20  $\beta^+$  decay is the emission of a positron that eventually finds an electron to annihilate, characteristically producing gammas in opposite directions.

The antielectron is often represented by the symbol  $e^+$ , but in beta decay it is written as  $\beta^+$  to indicate the antielectron was emitted in a nuclear decay. Antielectrons are the antimatter counterpart to electrons, being nearly identical, having the same mass, spin, and so on, but having a positive charge and an electron family number of -1. When a **positron** encounters an electron, there is a mutual annihilation in which all the mass of the antielectron-electron pair is converted into pure photon energy. (The reaction,  $e^+ + e^- \rightarrow \gamma + \gamma$ , conserves electron family number as well as all other conserved quantities.) If a nuclide  ${}^A_A X_N$  is known to  $\beta^+$  decay, then its  $\beta^+$  **decay equation** is

$${}^{A}_{Z}X_{N} \rightarrow {}^{A}_{Z-1}Y_{N+1} + \beta^{+} + \nu_{e} \ (\beta^{+} \text{ decay}),$$
 31.30

where Y is the nuclide having one less proton than X (to conserve charge) and  $\nu_e$  is the symbol for the **electron's neutrino**, which has an electron family number of +1. Since an antimatter member of the electron family (the  $\beta^+$ ) is created in the decay, a matter member of the family (here the  $\nu_e$ ) must also be created. Given, for example, that <sup>22</sup> Na  $\beta^+$  decays, you can write its full decay equation by first finding that Z = 11 for <sup>22</sup> Na, so that the daughter nuclide will have Z = 10, the atomic number for neon. Thus the  $\beta^+$  decay equation for <sup>22</sup> Na is

$${}^{22}_{11}\mathrm{Na}_{11} \to {}^{22}_{10}\mathrm{Ne}_{12} + \beta^+ + \nu_e.$$

In  $\beta^+$  decay, it is as if one of the protons in the parent nucleus decays into a neutron, a positron, and a neutrino. Protons do not do this outside of the nucleus, and so the decay is due to the complexities of the nuclear force. Note again that the total number of nucleons is constant in this and any other reaction. To find the energy emitted in  $\beta^+$  decay, you must again count the number of electrons in the neutral atoms, since atomic masses are used. The daughter has one less electron than the parent, and one electron mass is created in the decay. Thus, in  $\beta^+$  decay,

$$\Delta m = m(\text{parent}) - [m(\text{daughter}) + 2m_e], \qquad 31.32$$

since we use the masses of neutral atoms.

**Electron capture** is the third type of beta decay. Here, a nucleus captures an inner-shell electron and undergoes a nuclear reaction that has the same effect as  $\beta^+$  decay. Electron capture is sometimes denoted by the letters EC. We know that electrons cannot reside in the nucleus, but this is a nuclear reaction that consumes the electron and occurs spontaneously only when the products have less mass than the parent plus the electron. If a nuclide  ${}^A_Z X_N$  is known to undergo electron capture, then its **electron capture equation** is

$${}^{A}_{Z}X_{N} + e^{-} \rightarrow {}^{A}_{Z-1}Y_{N+1} + \nu_{e}$$
 (electron capture, or EC). 31.33

Any nuclide that can  $\beta^+$  decay can also undergo electron capture (and often does both). The same conservation laws are obeyed for EC as for  $\beta^+$  decay. It is good practice to confirm these for yourself.

All forms of beta decay occur because the parent nuclide is unstable and lies outside the region of stability in the chart of nuclides. Those nuclides that have relatively more neutrons than those in the region of stability will  $\beta^-$  decay to produce a daughter with fewer neutrons, producing a daughter nearer the region of stability. Similarly, those nuclides having relatively more protons than those in the region of stability will  $\beta^-$  decay or undergo electron capture to produce a daughter with fewer protons, nearer the region of stability.

## Gamma Decay

**Gamma decay** is the simplest form of nuclear decay—it is the emission of energetic photons by nuclei left in an excited state by some earlier process. Protons and neutrons in an excited nucleus are in higher orbitals, and they fall to lower levels by photon emission (analogous to electrons in excited atoms). Nuclear excited states have lifetimes typically of only about  $10^{-14}$  s, an

indication of the great strength of the forces pulling the nucleons to lower states. The  $\gamma$  decay equation is simply

$${}^{A}_{Z}X_{N}^{*} \rightarrow {}^{A}_{Z}X_{N} + \gamma_{1} + \gamma_{2} + \cdots (\gamma \text{ decay})$$

$$31.34$$

where the asterisk indicates the nucleus is in an excited state. There may be one or more  $\gamma$  s emitted, depending on how the nuclide de-excites. In radioactive decay,  $\gamma$  emission is common and is preceded by  $\gamma$  or  $\beta$  decay. For example, when  ${}^{60}$ Co  $\beta^-$  decays, it most often leaves the daughter nucleus in an excited state, written  ${}^{60}$ Ni\*. Then the nickel nucleus quickly  $\gamma$  decays by the emission of two penetrating  $\gamma$  s:

$${}^{60}\text{Ni}^* \to {}^{60}\text{Ni} + \gamma_1 + \gamma_2.$$
 31.35

These are called cobalt  $\gamma$  rays, although they come from nickel—they are used for cancer therapy, for example. It is again constructive to verify the conservation laws for gamma decay. Finally, since  $\gamma$  decay does not change the nuclide to another species, it is not prominently featured in charts of decay series, such as that in Figure 31.16.

There are other types of nuclear decay, but they occur less commonly than  $\alpha$ ,  $\beta$ , and  $\gamma$  decay. Spontaneous fission is the most important of the other forms of nuclear decay because of its applications in nuclear power and weapons. It is covered in the next chapter.

## **31.5 Half-Life and Activity**

Unstable nuclei decay. However, some nuclides decay faster than others. For example, radium and polonium, discovered by the Curies, decay faster than uranium. This means they have shorter lifetimes, producing a greater rate of decay. In this section we explore half-life and activity, the quantitative terms for lifetime and rate of decay.

## Half-Life

Why use a term like half-life rather than lifetime? The answer can be found by examining Figure 31.21, which shows how the number of radioactive nuclei in a sample decreases with time. The *time in which half of the original number of nuclei decay* is defined as the **half-life**,  $t_{1/2}$ . Half of the remaining nuclei decay in the next half-life. Further, half of that amount decays in the following half-life. Therefore, the number of radioactive nuclei decreases from N to N/2 in one half-life, then to N/4 in the next, and to N/8 in the next, and so on. If N is a large number, then *many* half-lives (not just two) pass before all of the nuclei decay. Nuclear decay is an example of a purely statistical process. A more precise definition of half-life is that *each nucleus has a 50% chance of living for a time equal to one half-life*  $t_{1/2}$ . Thus, if N is reasonably large, half of the original nuclei decay in a time of one half-life. If an individual nucleus makes it through that time, it still has a 50% chance of surviving through one more. The probability of decay is the same no matter when you start counting. This is like random coin flipping. The chance of heads is 50%, no matter what has happened before.



Figure 31.21 Radioactive decay reduces the number of radioactive nuclei over time. In one half-life  $t_{1/2}$ , the number decreases to half of its original value. Half of what remains decay in the next half-life, and half of those in the next, and so on. This is an exponential decay, as seen in the graph of the number of nuclei present as a function of time.

There is a tremendous range in the half-lives of various nuclides, from as short as  $10^{-23}$  s for the most unstable, to more than